













Part. & Energy conservation [Case: 1.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_p = 101$]
 Comparison with initial solution - log scale; total time and zoom over time







Part. & Energy conservation [Case: 1.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_p = 101$]
 Comparison with initial solution - linear scale; total time and zoom over time



Part. & Energy conservation [Case: I.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_p = 101$]
Comparison with previous time-sampled (τ_{out}) solution - log and linear scales







Particle conservation [Case: 1.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_p = 101$]
 Comparison with asymptotic solution (electrons and ions); total time and zoom over time







Profiles [Case: 1.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_p = 101$]

Time sampling: total simulation time/10



Profiles [Case: 1.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_\rho = 101$]





Comparison with asymptotic solution



● final calculation
● asymptotic



Profiles [Case: 1.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_\rho = 101$]

Time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (V a/D)| = 0.80 \text{ s}$







0.00
0.10
0.20
0.30
0.40
0.50
0.60
0.70





Profiles [Case: I.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_\rho = 101$]
 Spatial zoom over magnetic axis; time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (Va/D)| = 0.80 \text{ s}$



Profiles [Case: 1.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_\rho = 101$]
 Spatial zoom over edge; time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (V_a/D)| = 0.80 \text{ s}$





Profiles [Case: I.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_p = 101$]
 Time sampling: last 10 time slices



9.00
 9.10
 9.20
 9.30
 9.40
 9.50
 9.60
 9.70
 9.80
 9.90

Profiles [Case: 1.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_\rho = 101$]
 Spatial zoom over magnetic axis; time sampling: last 10 time slices



Profiles [Case: 1.1.5.b, Solver: 7, $D = 0.1 \text{ m}^2/\text{s}$, $v = 0.30 \text{ m/s}$, $\Delta t = 10.01$, $\tau = 1.0 \times 10^{-2} \text{ s}$, $N_\rho = 101$]
 Spatial zoom over edge; time sampling: last 10 time slices

