Part. & Energy conservation [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m}/\mathrm{s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Comparison with initial solution - log scale; total time and zoom over time









Part. & Energy conservation [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m}/\mathrm{s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Comparison with initial solution - linear scale; total time and zoom over time









Part. & Energy conservation [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m}/\mathrm{s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Comparison with previous time-sampled (τ_{out}) solution - log and linear scales

Particle conservation (electrons and ions)



Energy conservation (electrons and ions)







Particle conservation [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m}/\mathrm{s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Comparison with asymptotic solution (electrons and ions); total time and zoom over time









Profiles [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m/s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Time sampling: total simulation time/10











Profiles [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m}/\mathrm{s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Comparison with asymptotic solution









Profiles [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m/s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (Va/D)|$ =4.00 s









Profiles [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m/s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Spatial zoom over magnetic axis; time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (Va/D)|$ =4.00 s



Profiles [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m/s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Spatial zoom over edge; time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (Va/D)|$ =4.00 s









Profiles [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m/s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Time sampling: last 10 time slices









Profiles [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m/s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Spatial zoom over magnetic axis; time sampling: last 10 time slices









Profiles [Case: I.1.5, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = 0.00\,\mathrm{m/s}$, Δt = 50.00, $\tau = 1.0 \times 10^{-3}\,\mathrm{s}$, $N_\rho = 101$]

Spatial zoom over edge; time sampling: last 10 time slices